

## Package leaflet: information for the patient

### Procoralan 5 mg film-coated tablets Procoralan 7.5 mg film-coated tablets ivabradine

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

#### **Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

1. What Procoralan is and what it is used for
2. What you need to know before you take Procoralan
3. How to take Procoralan
4. Possible side effects
5. How to store Procoralan
6. Contents of the pack and other information

#### **1. What Procoralan is and what it is used for**

Procoralan (ivabradine) is a heart medicine used to treat:

- Symptomatic stable angina pectoris (which causes chest pain) in adult patients whose heart rate is over or equal to 70 beats per minute. It is used in adult patients who do not tolerate or cannot take heart medicines called beta-blockers. It is also used in combination with beta-blockers in adult patients whose condition is not fully controlled with a beta-blocker.
- Chronic heart failure in adult patients whose heart rate is over or equal to 75 beats per minute. It is used in combination with standard therapy, including beta-blocker therapy or when beta-blockers are contraindicated or not tolerated.

#### **About stable angina pectoris (usually referred to as “angina”):**

Stable angina is a heart disease which happens when the heart does not receive enough oxygen. It usually appears between 40 and 50 years of age. The most common symptom of angina is chest pain or discomfort. Angina is more likely to happen when the heart beats faster in situations such as exercise, emotion, exposure to the cold or after eating. This increase in heart rate can cause the chest pain in people who suffer from angina.

#### **About chronic heart failure :**

Chronic heart failure is a heart disease which happens when your heart cannot pump enough blood to the rest of your body. The most common symptoms of heart failure are breathlessness, fatigue, tiredness and ankle swelling.

#### **How does Procoralan work?**

Procoralan mainly works by reducing the heart rate by a few beats per minute. This lowers the heart's need for oxygen especially in the situations when an angina attack is more likely to happen. In this way Procoralan helps to control and reduce the number of angina attacks.

Furthermore as elevated heart rate adversely affects the heart functioning and vital prognosis in patients with chronic heart failure, the specific heart rate lowering action of ivabradine helps to improve the heart functioning and vital prognosis in these patients.

## **2. What you need to know before you take Procoralan**

### **Do not take Procoralan**

- if you are allergic to ivabradine or any of the other ingredients of this medicine (listed in section 6);
- if your resting heart rate before treatment is too slow (below 70 beats per minute);
- if you are suffering from cardiogenic shock (a heart condition treated in hospital);
- if you suffer from a heart rhythm disorder;
- if you are having a heart attack;
- if you suffer from very low blood pressure;
- if you suffer from unstable angina (a severe form in which chest pain occurs very frequently and with or without exertion);
- if you have heart failure which has recently become worse;
- if your heart beat is exclusively imposed by your pacemaker;
- if you suffer from severe liver problems;
- if you are already taking medicines for the treatment of fungal infections (such as ketoconazole, itraconazole), macrolide antibiotics (such as josamycin, clarithromycin, telithromycin or erythromycin given orally), medicines to treat HIV infections (such as nelfinavir, ritonavir) or nefazodone (medicine to treat depression) or diltiazem, verapamil (used for high blood pressure or angina pectoris);
- if you are a woman able to have children and not using reliable contraception;
- if you are pregnant or trying to become pregnant;
- if you are breast-feeding.

### **Warnings and precautions**

Talk to your doctor or pharmacist before taking Procoralan

- if you suffer from heart rhythm disorders (such as irregular heartbeat, palpitation, increase in chest pain) or sustained atrial fibrillation (a type of irregular heartbeat), or an abnormality of electrocardiogram (ECG) called 'long QT syndrome',
- if you have symptoms such as tiredness, dizziness or shortness of breath (this could mean that your heart is slowing down too much),
- if you suffer from symptoms of atrial fibrillation (pulse rate at rest unusually high (over 110 beats per minute) or irregular, without any apparent reason, making it difficult to measure),
- if you have had a recent stroke (cerebral attack),
- if you suffer from mild to moderate low blood pressure,
- if you suffer from uncontrolled blood pressure, especially after a change in your antihypertensive treatment,
- if you suffer from severe heart failure or heart failure with abnormality of ECG called 'bundle branch block',
- if you suffer from chronic eye retinal disease,
- if you suffer from moderate liver problems,
- if you suffer from severe renal problems.

If any of the above applies to you, talk straight away to your doctor before or while taking Procoralan.

### **Children**

Procoralan is not intended for use in children and adolescents younger than 18 years.

### **Other medicines and Procoralan**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Make sure to tell your doctor if you are taking any of the following medicines, as a dose adjustment of Procoralan or monitoring should be required:

- fluconazole (an antifungal medicine)
- rifampicin (an antibiotic)
- barbiturates (for difficult sleeping or epilepsy)
- phenytoin (for epilepsy)
- *Hypericum perforatum* or St John's Wort (herbal treatment for depression)
- QT prolonging medicines to treat either heart rhythm disorders or other conditions :
  - quinidine, disopyramide, ibutilide, sotalol, amiodarone (to treat heart rhythm disorders)
  - bepridil (to treat angina pectoris)
  - certain types of medicines to treat anxiety, schizophrenia or other psychoses (such as pimozone, ziprasidone, sertindole)
  - anti-malarial medicines (such as mefloquine or halofantrine)
  - intravenous erythromycin (an antibiotic)
  - pentamidine (an antiparasitic medicine)
  - cisapride (against the gastro-oesophageal reflux)
- Some types of diuretics which may cause decrease in blood potassium level, such as furosemide, hydrochlorothiazide, indapamide (used to treat oedema, high blood pressure).

### **Procoralan with food and drink**

Avoid grapefruit juice during treatment with Procoralan.

### **Pregnancy and breast-feeding**

Do not take Procoralan if you are pregnant or are planning to have a baby (see "Do not take Procoralan").

If you are pregnant and have taken Procoralan, talk to your doctor.

Do not take Procoralan if you are able to become pregnant unless you use reliable contraceptive measures (see "Do not take Procoralan").

Do not take Procoralan if you are breast-feeding (see "Do not take Procoralan"). Talk to your doctor if you are breast-feeding or intending to breast-feed as breastfeeding should be discontinued if you take Procoralan.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

### **Driving and using machines**

Procoralan may cause temporary luminous visual phenomena (a temporary brightness in the field of vision, see "Possible side effects"). If this happens to you, be careful when driving or using machines at times when there could be sudden changes in light intensity, especially when driving at night.

### **Procoralan contains lactose**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

## **3. How to take Procoralan**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Procoralan should be taken during meals.

### If you are being treated for stable angina pectoris

The starting dose should not exceed one tablet of Procoralan 5 mg twice daily. If you still have angina symptoms and if you have tolerated the 5 mg twice daily dose well, the dose may be increased. The maintenance dose should not exceed 7.5 mg twice daily. Your doctor will prescribe the right dose for you. The usual dose is one tablet in the morning and one tablet in the evening. In some cases (e.g. if

you are elderly), your doctor may prescribe half the dose i.e., one half 5 mg tablet of Procoralan 5 mg (corresponding to 2.5 mg ivabradine) in the morning and one half 5 mg tablet in the evening.

**If you are being treated for chronic heart failure**

The usual recommended starting dose is one tablet of Procoralan 5 mg twice daily increasing if necessary to one tablet of Procoralan 7.5 mg twice daily. Your doctor will decide the right dose for you. The usual dose is one tablet in the morning and one tablet in the evening. In some cases (e.g. if you are elderly), your doctor may prescribe half the dose i.e., one half 5 mg tablet of Procoralan 5 mg (corresponding to 2.5 mg ivabradine) in the morning and one half 5 mg tablet in the evening.

**If you take more Procoralan than you should:**

A large dose of Procoralan could make you feel breathless or tired because your heart slows down too much. If this happens, contact your doctor immediately.

**If you forget to take Procoralan:**

If you forget to take a dose of Procoralan, take the next dose at the usual time. Do not take a double dose to make up for the forgotten dose.

The calendar printed on the blister containing the tablets should help you remember when you last took a tablet of Procoralan.

**If you stop taking Procoralan:**

As the treatment for angina or chronic heart failure is usually life-long, you should discuss with your doctor before stopping this medicinal product.

If you think that the effect of Procoralan is too strong or too weak, talk to your doctor or pharmacist.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The frequency of possible side effects listed below is defined using the following convention:

very common: may affect more than 1 in 10 people

common: may affect up to 1 in 10 people

uncommon: may affect up to 1 in 100 people

rare: may affect up to 1 in 1,000 people

very rare: may affect up to 1 in 10,000 people

not known: frequency cannot be estimated from the available data

The most common adverse reactions with this medicine are dose dependent and related to its mode of action:

*Very common:*

Luminous visual phenomena (brief moments of increased brightness, most often caused by sudden changes in light intensity). They can also be described as a halo, coloured flashes, image decomposition or multiple images. They generally occur within the first two months of treatment after which they may occur repeatedly and resolve during or after treatment

*Common:*

Modification in the heart functioning (the symptoms are a slowing down of the heart rate). They particularly occur within the first 2 to 3 months of treatment initiation.

Other side effects have also been reported:

*Common:*

Irregular rapid contraction of the heart, abnormal perception of heartbeat, uncontrolled blood pressure, headache, dizziness and blurred vision (cloudy vision).

*Uncommon:*

Palpitations and cardiac extra beats, feeling sick (nausea), constipation, diarrhoea, abdominal pain, spinning sensation (vertigo), difficulty breathing (dyspnoea), muscle cramps, changes in laboratory parameters : high blood levels of uric acid, an excess of eosinophils (a type of white blood cell) and elevated creatinine in blood (a breakdown product of muscle), skin rash, angioedema (such as swollen face, tongue or throat, difficulty in breathing or swallowing), low blood pressure, fainting, feeling of tiredness, feeling of weakness, abnormal ECG heart tracing, double vision, impaired vision.

*Rare:*

Urticaria, itching, skin reddening, feeling unwell.

*Very rare:*

Irregular heart beats.

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via **the national reporting system listed in Appendix V**. By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Procoralan**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after 'EXP'. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.


Do not throw away any medicine via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.


## **6. Content of the pack and other information**

### **What Procoralan contains**

- The active substance is ivabradine (as hydrochloride).  
Procoralan 5 mg: one film-coated tablet contains 5 mg ivabradine (equivalent to 5.390 mg ivabradine as hydrochloride).  
Procoralan 7.5 mg: one film-coated tablet contains 7.5 mg ivabradine (equivalent to 8.085 mg ivabradine as hydrochloride).
- The other ingredients in the tablet core are: lactose monohydrate, magnesium stearate (E 470 B), maize starch, maltodextrin, colloidal anhydrous silica (E 551), and in the tablet coating: hypromellose (E 464), titanium dioxide (E 171), macrogol 6000, glycerol (E 422), magnesium stearate (E 470 B), yellow iron oxide (E 172), red iron oxide (E 172).

### **What Procoralan looks like and contents of the pack**

Procoralan 5 mg tablets are salmon-coloured, oblong film-coated tablets scored on both sides, engraved with "5" on one face and  on the other.

Procoralan 7.5 mg tablets are salmon-coloured, triangular, film-coated tablets engraved with "7.5" on one face and  on the other.

The tablets are available in calendar packs (Aluminium/PVC blisters) of 14, 28, 56, 84, 98, 100 or 112 tablets.

Not all pack sizes may be marketed.

**Marketing Authorisation Holder and Manufacturer:**

Marketing Authorisation Holder:

Les Laboratoires Servier  
50, rue Carnot  
92284 Suresnes cedex - France

Manufacturer:

Les Laboratoires Servier Industrie  
905 route de Saran  
45520 Gidy – France

Servier (Ireland) Industries Ltd  
Gorey Road  
Arklow - Co. Wicklow – Ireland

Przedsiębiorstwo Farmaceutyczne ANPHARM S.A.  
ul. Annopol 6B – 03-236 Warszawa – Poland

and

Laboratorios Servier, S.L.  
Avda. de los Madroños, 33  
28043 Madrid  
Spain

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

**België/Belgique/Belgien**

S.A. Servier Benelux N.V.  
Tél/Tel: +32 (0)2 529 43 11

**България**

Сервие Медикал ЕООД  
Тел.: +359 2 921 57 00

**Česká republika**

Servier s.r.o.  
Tel: +420 222 118 111

**Danmark**

Servier Danmark A/S  
Tlf: +45 36 44 22 60

**Deutschland**

Servier Deutschland GmbH  
Tel: +49 (0)89 57095 01

**Lietuva**

UAB "SERVIER PHARMA"  
Tel: +370 (5) 2 63 86 28

**Luxembourg/Luxemburg**

S.A. Servier Benelux N.V.  
Tél/Tel: +32 (0)2 529 43 11

**Magyarország**

Servier Hungaria Kft.  
Tel.: + 36 1 238 77 99

**Malta**

GALEPHARMA Ltd  
Tel: +(356) 21 247 082

**Nederland**

Servier Nederland Farma B.V.  
Tel: +31 (0)71 5246700

**Eesti**

Servier Laboratories OÜ  
Tel: +372 664 5040

**Ελλάδα**

ΣΕΡΒΙΕ ΕΛΛΑΣ ΦΑΡΜΑΚΕΥΤΙΚΗ ΕΠΕ  
Τηλ: +30 210 939 1000

**España**

Laboratorios Servier S.L.  
Tel: +34 91 748 96 30

**France**

Les Laboratoires Servier  
Tél: +33 (0)1 55 72 60 00

**Hrvatska**

Servier Pharma, d. o. o.  
Tel.: +385 (0)1 3016 222

**Ireland**

Servier Laboratories (Ireland) Ltd.  
Tel: +353 (0)1 663 8110

**Ísland**

Servier Laboratories  
C/o Icepharma hf  
Sími: +354 540 8000

**Italia**

Servier Italia S.p.A.  
Tel: +39 06 669081

**Κύπρος**

C.A. Papaellinas Ltd.  
Τηλ: +357 22741741

**Latvija**

SIA Servier Latvia  
Tel: +371 6750 2039

**Norge**

Servier Danmark A/S  
Tlf: +45 36 44 22 60

**Österreich**

Servier Austria GmbH  
Tel: +43 (1) 524 39 99

**Polska**

Servier Polska SP. Z O.O.  
Tel.: +48 (0) 22 594 90 00

**Portugal**

Servier Portugal, Lda  
Tel: +351 21 312 20 00

**România**

Servier Pharma SRL  
Tel: +4 021 528 52 80

**Slovenija**

Servier Pharma d.o.o.  
Tel: +386 (0)1 563 48 11

**Slovenská republika**

Servier Slovensko spol. s r.o.  
Tel: +421 (0) 2 5920 41 11

**Suomi/Finland**

Servier Finland Oy  
P. /Tel: +358 (0)9 279 80 80

**Sverige**

Servier Sverige AB  
Tel: +46(8)5 225 08 00

**United Kingdom**

Servier Laboratories Ltd  
Tel: +44 (0)1753 666409

**This leaflet was last revised in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>